

**Significance of the non-2-spinon part of $S(q, \omega)$ in the
one-dimensional $S = 1/2$ Heisenberg antiferromagnet at zero
temperature**

Andreas Fledderjohann and Karl-Heinz Mütter

Physics Department, University of Wuppertal, 42097 Wuppertal, Germany

Michael Karbach and Gerhard Müller

Department of Physics, The University of Rhode Island, Kingston RI 02881-0817

(February 1, 2008)

Abstract

The impact of the non-2-spinon excitations of the one-dimensional $S = 1/2$ Heisenberg antiferromagnet on the integrated intensity, the susceptibility, the frequency moments, and the Euclidian time representation of the $T = 0$ dynamic spin structure factor $S(q, \omega)$ is studied on the basis of finite-size data for chains with up to $N = 28$ sites.

PACS number: 75.10 -b, 75.10.Jm, 75.40.Gb

I. INTRODUCTION

The spectral weight in the dynamic structure factor

$$S(q, \omega) \equiv \frac{1}{N} \sum_{l,n} e^{iqn} \int_{-\infty}^{+\infty} dt e^{i\omega t} \langle S_l^z(t) S_{l+n}^z \rangle \quad (1)$$

of the one-dimensional (1D) $S = 1/2$ Heisenberg antiferromagnet

$$H = J \sum_{l=1}^N \mathbf{S}_l \cdot \mathbf{S}_{l+1} \quad (2)$$

at $T = 0$ has long been known to come predominantly from the 2-spinon excitations,¹ which form a continuum between the lower and upper boundaries (with energies in units of J)^{2,1}

$$\omega_L(q) = \frac{\pi}{2} \sin q, \quad \omega_U(q) = \pi \sin \frac{q}{2}. \quad (3)$$

However, the exact 2-spinon part of $S(q, \omega)$, here named $S^{(2)}(q, \omega)$, was determined only very recently,^{3,4} based on new advances in quantum groups.⁵

The exact expression of $S^{(2)}(q, \omega)$, which had a fairly complex structure in its original form,^{3,5} was simplified considerably in Ref. 4, evaluated numerically, and plotted versus q and ω . The spectral-weight distribution was found to diverge at $\omega_L(q)$,

$$S^{(2)}(q, \omega) \sim \frac{1}{\sqrt{\omega - \omega_L(q)}} \sqrt{\ln \frac{1}{\omega - \omega_L(q)}}, \quad (4)$$

and to vanish in a square-root cusp at $\omega_U(q)$,

$$S^{(2)}(q, \omega) \sim \sqrt{\omega_U(q) - \omega}. \quad (5)$$

These singularities differ from those of the approximate expression

$$S^{(a)}(q, \omega) = \frac{\Theta(\omega - \omega_L(q)) \Theta(\omega_U(q) - \omega)}{\sqrt{\omega^2 - \omega_L(q)^2}} \quad (6)$$

for the 2-spinon dynamic structure factor, which had been inferred from finite- N data, sum rules, and Bethe-ansatz calculations.¹ Otherwise, the line shapes of $S^{(2)}(q, \omega)$ and $S^{(a)}(q, \omega)$ do not differ all that much.⁴

The 2-spinon excitations were shown to account for 82.18% of the total intensity in $S(q, \omega)$. Where in (q, ω) -space is the remaining spectral weight, from which classes of Bethe-ansatz solutions does it originate, and how does it affect various quantities that can be derived from the dynamic structure factor?

A study of the 4-spinon contribution to $S(q, \omega)$ along the lines of Refs. 5,3,4 will shed light on the first two questions.⁶ In answer to the third question, we investigate here the effects of the non-2-spinon excitations on four quantities which are related to $S(q, \omega)$ and which can be computed with high precision from finite- N data for the ground-state wave function.

II. INTEGRATED INTENSITY

The integrated intensity of $S(q, \omega)$, i.e. the static spin structure factor

$$I(q) \equiv \int_0^\infty \frac{d\omega}{2\pi} S(q, \omega), \quad (7)$$

has been determined with high precision for wave numbers $q \leq 13\pi/14$ from finite- N data of cyclic chains with $N \leq 28$ sites.⁷ This result is plotted in Fig. 1 for comparison with the exact 2-spinon integrated intensity $I^{(2)}(q)$ calculated from $S^{(2)}(q, \omega)$ via (7) and the integrated intensity¹

$$I^{(a)}(q) = \frac{1}{2\pi} \ln \frac{1 + \sin(q/2)}{\cos(q/2)} \quad (8)$$

obtained from the approximate result $S^{(a)}(q, \omega)$.

The finite- N data indicate that $I(q)$ increases linearly from zero for small q and diverges logarithmically at $q = \pi$. The initial rise of the finite- N data, $I(q) \rightarrow 0.271q/\pi$,⁸ is significantly steeper than that of the 2-spinon contribution, $I^{(2)}(q) \rightarrow 0.237q/\pi$ and that of the approximate result, $I^{(a)}(q) \rightarrow 0.25q/\pi$. Hence the integrated intensity of the non-2-spinon part of $S(q, \omega)$ increases linearly in q too.

The function $I^{(a)}(q)$ approximates the 2-spinon integrated intensity $I^{(2)}(q)$ quite well for $q/\pi \lesssim 0.6$. At larger q , this is no longer the case. The divergence predicted by expression

(8), $I^{(a)} \sim -\ln(1 - q/\pi)$, is weaker than the divergence of the exact 2-spinon result, $I^{(2)}(q) \sim [-\ln(1 - q/\pi)]^{3/2}$. The inset of Fig. 1 shows the relative non-2-spinon integrated intensity, $\Delta I^{(2)}(q) = 1 - I^{(2)}(q)/I(q)$, and the relative deviation, $\Delta I^{(a)}(q) = 1 - I^{(a)}(q)/I(q)$, of the approximate result (8). If it can be assumed that the leading singularity of $I(q)$ at $q = \pi$ is produced entirely by the 2-spinon part of $S(q, \omega)$, then the function $\Delta I^{(2)}(q)$ must approach zero as $q \rightarrow \pi$. The dashed line in the inset does not rule out that this assumption is correct. Whether or not the 4-spinon excitations contribute to the leading singularity in $I(q)$ remains to be seen.⁶

It is interesting to compare these results with the exact integrated intensity of the Haldane-Shastry model,⁹ $I^{(HS)}(q) = -(1/4)\ln(1 - q/\pi)$, where non-2-spinon excitations have zero spectral weight in $S(q, \omega)$. It turns out that for $q \lesssim 13\pi/14$, $I^{(HS)}(q)$ is a much better approximation of $I(q)$ than $I^{(2)}(q)$ is.⁷

III. SUSCEPTIBILITY

The q -dependent susceptibility at $T = 0$ is related to $S(q, \omega)$ via the sum rule¹⁰

$$\chi(q) \equiv \frac{1}{\pi} \int_0^\infty \frac{d\omega}{\omega} S(q, \omega). \quad (9)$$

This quantity, which has been determined with considerable accuracy from finite- N data, is plotted in Fig. 2 for comparison with the exact 2-spinon susceptibility $\chi^{(2)}(q)$ calculated⁴ from $S^{(2)}(q, \omega)$ via (9) and the approximate result¹

$$\chi^{(a)}(q) = \frac{1}{\pi^2} \frac{q}{\sin q} \quad (10)$$

inferred from (6).

The normalization of $S^{(a)}(q, \omega)$ was chosen such that the exact value of the direct susceptibility,¹¹ $\chi(0) = 1/\pi^2$, is correctly reproduced. With increasing q , $\chi^{(a)}(q)$ deviates in a downward direction from $\chi(q)$. Its divergence at $q = \pi$, $\chi^{(a)}(q) \sim (\pi - q)^{-1}$, is slightly weaker than that of the exact 2-spinon susceptibility,⁴ $\chi^{(2)}(q) \sim \sqrt{-\ln(\pi - q)}/(\pi - q)$, which, presumably coincides with the leading singularity of $\chi(q)$.

The contribution of the non-2-spinon spectral weight of $S(q, \omega)$ to $\chi(q)$ in the limit $q \rightarrow 0$ is small but non-negligible as indicated by the result $\pi^2 \chi^{(2)}(0) = 0.9500\dots$, which falls some 5% short of the exact direct susceptibility. The relative non-2-spinon contribution to the susceptibility, $\Delta\chi^{(2)}(q) = 1 - \chi^{(2)}(q)/\chi(q)$, stays smaller for all q -values than the relative non-2-spinon integrated intensity $\Delta I^{(2)}(q)$ (see inset). This indicates that the non-2-spinon spectral weight is located predominately above the 2-spinon continuum.

IV. FREQUENCY MOMENTS

Yet a different way to assess the non-2-spinon part of $S(q, \omega)$ employs the frequency moments

$$K_n(q) \equiv \int_0^\infty \frac{d\omega}{2\pi} \omega^n S(q, \omega), \quad n = 1, 2, \dots, \quad (11)$$

which are related, via sum rules,¹² to short-range multi-spin correlations in the ground state. For $n = 1$ we know the exact results,¹⁰

$$K_1(q) = \frac{2E_0}{3N}(1 - \cos q), \quad (12)$$

where $E_0 = -N(\ln 2 - 1/4)$ is the ground-state energy. For $n = 2, 3, 4, 5$ high-precision results have been calculated from finite- N data for the associated ground-state expectation values.⁸ The moments $K_n^{(2)}(q)$ of the exact 2-spinon dynamic structure factor $S^{(2)}(q, \omega)$ have been determined in Ref. 4 and the moments $K_n^{(a)}(q)$ of $S^{(a)}(q, \omega)$ in Ref. 12.

For $n = 1$, both $K_1^{(2)}(q)$ and $K_1^{(a)}(q)$ reproduce the q -dependence of the exact sum rule (12) correctly, but the prefactors are smaller,

$$\frac{K_1^{(2)}(q)}{K_1(q)} = 0.8039\dots, \quad \frac{K_1^{(a)}(q)}{K_1(q)} = 0.8462\dots, \quad (13)$$

which again reflects the missing spectral weight of the non-2-spinon excitations. The q -dependence of the moment ratios

$$R_n(q) \equiv \frac{K_n(q)}{K_1(q)}, \quad \text{for } n = 2, 3, 4, 5, \quad (14)$$

of the full dynamic structure factor as inferred from finite- N data are shown in Fig. 3 along with the corresponding moment ratios $R_n^{(2)}(q)$ of $S^{(2)}(q, \omega)$ and the moment ratios $R_n^{(a)}(q)$ of $S^{(a)}(q, \omega)$. The most striking observation is that $R_n(q)$ approaches a nonzero value as $q \rightarrow 0$, whereas the exact and the approximate 2-spinon moment ratios both go to zero: $R_n^{(2)}(q) \sim R_n^{(a)}(q) \sim q^{n-1}$. This means that for long wavelengths the frequency moments $K_n(q)$, $n \geq 2$, are dominated by non-2-spinon excitations, which are necessarily located above the narrow 2-spinon band. In other words, the 2-spinon dynamic structure factor $S^{(2)}(q, \omega)$ does not contribute to the leading $O(q^2)$ term of $K_n(q)$ for $n \geq 2$.

At larger wave numbers, the impact of the non-2-spinon excitations on the moment ratios is more modest but still significant. Here the deviation of $R_n^{(2)}(q)$ from $R_n(q)$ is almost q -independent, and it grows with increasing n . This again indicates that the non-2-spinon spectral weight comes for the most part from higher frequencies than the 2-spinon spectral weight.

The moment ratios $R_n^{(a)}(q)$ agree very well with $R_n^{(2)}(q)$ at small q , but then deviate upwardly. For $q \gtrsim \pi/3$, they rise even above the ratios $R_n(q)$. This discrepancy, which becomes more conspicuous with increasing n , reflects the fact that $S^{(a)}(q, \omega)$ underestimates the spectral weight near the lower continuum boundary $\omega_L(q)$ and overestimates the spectral weight near the upper boundary $\omega_U(q)$.

V. EUCLIDIAN TIME REPRESENTATION

In order to study the significance of the non-2-spinon excitations in a dynamical quantity by the same kind of comparison, it is useful to consider the Laplace transform of the dynamic structure factor,¹³

$$\tilde{S}(q, \tau) \equiv \int_0^\infty \frac{d\omega}{2\pi} e^{-\omega\tau} S(q, \omega), \quad (15)$$

which can be interpreted as a Euclidian time representation of $S(q, \omega)$. For $\tau = 0$, this is the integrated intensity (7). From finite- N data for $S(q, \omega)$ as obtained via recursion method

for systems with $N \leq 28$ sites,¹³ this quantity can be accurately extrapolated to $N \rightarrow \infty$ if $q \neq \pi$. For the graphical representation, it is convenient to plot the function

$$\rho(q, t) \equiv \frac{\tilde{S}(q, 0)}{\tilde{S}(q, \tau)} - 1, \quad t = \sqrt{\omega_1(p)}\tau \exp[\omega_1(p)\tau], \quad (16)$$

which was used for the finite- N extrapolation, instead of $\tilde{S}(q, \tau)$ itself. The resulting curves, shown as solid lines in Fig. 4 for several q -values, rise from zero with zero initial slope and then become almost linear in t with a q -dependent slope.

If the threshold singularity were a square-root divergence as predicted by (6), then the asymptotic growth of $\rho(q, t)$ would be exactly linear. The logarithmic correction in the exact 2-spinon threshold singularity (4), however, leads to a slight modification of the asymptotic growth of $\rho(q, t)$,

$$\rho^{(2)}(q, t) \xrightarrow{t \rightarrow \infty} \propto \frac{t}{\ln \ln t}. \quad (17)$$

The dashed lines in Fig. 4 show the function $\rho^{(2)}(q, t)$ as inferred from the exact 2-spinon dynamic structure factor. The discrepancies are fairly small over the range of t shown. The deviation $\Delta^{(2)}(q, t) = \rho^{(2)}(q, t) - \rho(q, t)$ is shown in the inset. Not surprisingly, the function $\rho^{(a)}(q, t)$ inferred from (6) deviates more strongly from $\rho(q, t)$. This comparison was already made in Ref. 13.

ACKNOWLEDGMENTS

The work at URI was supported by NSF Grant DMR-93-12252, and by the Max Kade Foundation.

REFERENCES

- ¹ G. Müller, H. Thomas, G. Beck, and J.C. Bonner, Phys. Rev. B **24**, 1429 (1981);
- ² J. des Cloizeaux and J.J. Pearson, Phys. Rev. **128**, 2131 (1962); T. Yamada, Prog. Theor. Phys. **41**, 880 (1969).
- ³ A.H. Bougourzi, M. Couture, and M. Kacir, Preprint ITP-SB-96-21, Stony Brook, q-alg/9604019
- ⁴ M. Karbach, G. Müller, and A.H. Bougourzi, Preprint ITP-SB-96-27, Stony Brook, cond-mat/9606068.
- ⁵ M. Jimbo and T. Miwa, *Algebraic Analysis of Solvable Lattice Models*, (American Mathematical Society, CBMS, 1995).
- ⁶ A.H. Bougourzi, M. Karbach, and G. Müller, (unpublished)
- ⁷ M. Karbach and K.-H. Mütter, Z. Phys. B **90**, 83 (1993); M. Karbach, K.-H. Mütter, and M. Schmidt, Phys. Rev. B **50**, 9281 (1994).
- ⁸ A. Fledderjohann, M. Karbach, and K.-H. Mütter, Phys. Rev. B **53**, 11543 (1996).
- ⁹ F.D.M. Haldane, Phys. Rev. Lett. **60**, 635 (1988), **60**, 1866; B.M. Shastry, Phys. Rev. Lett. **60**, 639 (1988); F.D.M. Haldane, M.R. Zirnbauer, Phys. Rev. Lett. **71**, 4055 (1993); F.D.M. Haldane, J.C. Talstra, Phys. Rev. B **50**, 6889 (1994).
- ¹⁰ P.C. Hohenberg and W.F. Brinkmann Phys. Rev. B **10**, 128 (1974);
- ¹¹ R.B. Griffiths, Phys. Rev. **133**, A768 (1964).
- ¹² G. Müller, Phys. Rev. B **26**, 1311 (1982).
- ¹³ A. Fledderjohann, M. Karbach, K.-H. Mütter, and P. Wielath, J. Phys.: Condens. Matter **7**, 8993 (1995).

FIGURES

FIG. 1. Integrated intensities (7) in comparison: $I(q)$ is represented by finite- N data for $N = 6, 8, \dots, 28$. $I^{(2)}(q)$ is the exact 2-spinon result. $I^{(a)}(q)$ is the approximate result (8). The inset shows $\Delta I^{(2)}(q) = 1 - I^{(2)}(q)/I(q)$ and $\Delta I^{(a)}(q) = 1 - I^{(a)}(q)/I(q)$.

FIG. 2. The q -dependent susceptibility (9) in comparison: $\chi(q)$ is represented by finite- N data for $N = 6, 8, \dots, 28$. $\chi^{(2)}(q)$ is the exact 2-spinon result. $\chi^{(a)}(q)$ is the approximate result (10). The inset shows the relative non-2-spinon integrated intensity $\Delta\chi^{(2)}(q) = 1 - \chi^{(2)}(q)/\chi(q)$ and $\Delta\chi^{(a)}(q) = 1 - \chi^{(a)}(q)/\chi(q)$.

FIG. 3. Ratios of frequency moments (14). For $n = 2, 3, 4, 5$, the $R_n(q)$ represent finite- N data for $N = 6, \dots, 28$. The $R_n^{(2)}(q)$ are exact 2-spinon results and the $R_n^{(a)}(q)$ are the moment ratios for (6).

FIG. 4. The function (16) for $q = \pi/4, \pi/3, \pi/2, 2\pi/3, 3\pi/4$. (bottom to top) The solid lines represent extrapolated finite- N data and the dashed lines the exact 2-spinon-part of that function. The inset shows the non-2-spinon part $\Delta^{(2)}(q, t)$ of the same function.




FIG.1




FIG.2





FIG.3
(PART 1)





FIG.3
(PART 2)




FIG.4